

Tobacco Use

Table 6. Percent (standard error) of women who reported smoking cigarettes during the 3 months before pregnancy. Ques. #28

Maternal Characteristics	percent (s.e.)
Total	24.2(1.2)
Race	
Black	10.4(1.3)
White	33.9(1.6)
Age	
Less than 20	25.7(3.1)
20-29	27.1(1.5)
30+	16.9(1.9)
Education	
Less than High School	32.2(2.8)
High School	30.4(2.0)
More than High School	14.6(1.4)
Marital Status	
Married	22.4(1.5)
Other	25.6(1.8)
Birthweight	
LBW (<2500 grams)	22.4(3.2)
NBW (>= 2500 grams)	24.4(1.2)
Medicaid Status*	
Medicaid	29.4(1.7)
Other	17.5(1.5)

Table 7. Percent (standard error) of women who reported smoking cigarettes during the last trimester of pregnancy, according to sociodemographic characteristics. Ques. #29

Maternal Characteristics	percent (s.e.)
Total	12.8(0.9)
Race	
Black	5.4(1.0)
White	17.9(1.3)
Age	
Less than 20	14.2(2.4)
20-29	13.6(1.2)
30+	10.5(1.6)
Education	
Less than High School	22.1(2.5)
High School	14.9(1.5)
More than High School	5.9(0.9)
Marital Status	
Married	10.8(1.1)
Other	14.4(1.4)
Birthweight	
LBW (<2500 grams)	14.3(2.7)
NBW (>= 2500 grams)	12.7(1.0)
Medicaid Status*	
Medicaid	17.6(1.4)
Other	6.7(1.0)

Table 8. Percent (standard error) of women who reported smoking cigarettes during the 3-6 months after delivery, according to sociodemographic characteristics. Ques #30

Maternal Characteristics	percent (s.e.)
Total	20.5(1.1)
Race	
Black	10.0(1.3)
White	27.8(1.5)
Age	
Less than 20	24.4(3.0)
20-29	22.1(1.4)
30+	14.4(1.8)
Education	
Less than High School	30.6(2.7)
High School	24.7(1.9)
More than High School	11.2(1.2)
Marital Status	
Married	18.4(1.3)
Other	22.2(1.7)
Birthweight	
LBW (<2500 grams)	21.8(3.1)
NBW (>= 2500 grams)	20.3(1.1)
Medicaid Status*	
Medicaid	26.3(1.6)
Other	13.0(1.3)

* A mother is considered "on Medicaid" if she was enrolled in Medicaid during any of the following time periods: just before pregnancy, during prenatal care or during delivery.